

Joanna FOURQUET¹, Céline NOIROT², Christophe KLOPP²,
Philippe PINTON³, Sylvie COMBES¹, Claire HOEDE², Géraldine PASCAL¹

¹ GenPhySE, Université de Toulouse, INRAE, ENVT, F-31326, Castanet Tolosan, France

² INRAE, UR875 MIAT, PF Bioinfo GenoToul, F-31326, Castanet-Tolosan, France

³ Toxalim, Université de Toulouse, INRAE, ENVT, INP-Purpan, UPS, F-31027 Toulouse, France

Corresponding Author: geraldine.pascal@inrae.fr



ExpoMycopig project and JF are funded by France Futur Elevage

We are grateful to the Genotoul bioinformatics platform Toulouse Occitanie (Bioinfo Genotoul, doi: 10.15454/1.5572369328961167E12) for providing computing and storage resources

Acknowledgements

What bioinformatics solution exists to process whole metagenome shotgun data?

nf-core/mag

- Incomplete
 - ✓ Metagenome assembly
 - ✓ Taxonomic affiliation of reads and bins
 - ✗ Taxonomic affiliation of contigs
 - ✗ Functional annotation
- Easy of use
 - ✓ Automated with **nextflow** [1]
- Highly reproducible  [2]
 - ✓ Comes with a Singularity container
- Modular
 - ✓ Different parameters
 - ✓ Possibility to skip certain steps
- Available
 - ✓ <https://github.com/nf-core/mag>

ATLAS [3]

- Incomplete
 - ✓ Metagenome assembly
 - ✗ Taxonomic affiliation of reads and contigs
 - ✓ Taxonomic affiliation of bins
 - ✓ Functional annotation
- Issues during use
 - ✓ Automated with Snakemake [4]
- Reproducible
 - ✓ Installation and run with **CONDA**
- Modular
 - ✓ Different parameters
 - ✓ Possibility to skip certain steps
- Available
 - ✓ <https://github.com/metagenome-atlas/atlas>


Our solution: metagWGS

- Complete
 - ✓ Metagenome assembly
 - ✓ Taxonomic affiliation of reads, contigs and bins
 - ✓ Functional annotation
- Easy of use
 - ✓ Automated with **nextflow**
- Highly reproducible
 - ✓ Comes with a Singularity container  [2]
- Modular
 - ✓ Different parameters
 - ✓ Possibility to skip certain steps
- Available
 - ✓ <https://forgemia.inra.fr/genotoul-bioinfo/metagwgs>

metagWGS: preprocessing, assembly and annotation


Raw data

- Adapter sequence (entire or truncated)
- High quality read
- Low quality read
- Host read



Cleaning

- cutadapt [5]
- sickle [6]
- bwa mem [7]




Quality control

- FastQC [10]
- FastQC [10]


Assembly

metaspades [11] or megahit [12]



Reads deduplication

bwa mem, samtools markdup [13]




Assembly filter

Filter_contig_per_cpm.py




Taxonomic affiliation of reads

- kaiju [8]
- kronaTools [9]



Annotation of genes

prokka [14]



Possible skipping

metagWGS: quantification and affiliation


Clustering of genes

cd-hit-est [15]



Quantification of reads on genes

bwa mem, featureCounts [16]





	S1	S2
Gene 1	5+	1+		
Gene 2	2+	0+		
...				

Taxonomic affiliation of contigs

diamond [17], aln2taxaffi.py

#contig	consensus taxid	consensus lineage	S1	S2	S3
S1_1	1263	cellular organisms; Bacteria; Terrabacteria group; Firmicutes; Clostridia; Clostridiales			
S1_10	84108	cellular organisms; Bacteria; Terrabacteria group; Actinobacteria; Coriobacteriia			

Input from part 1



Annotation of genes

Reads deduplication

Assembly filter

Possible skipping


Binning of contigs

bowtie2 [18], metabat2 [19]



Taxonomic affiliation of bins







BAT [20]



nf-core/mag

nf-core/mag


MultiQC^[21] graphic outputs of metagWGS



Output tables and other graphs of metagWGS

Taxonomic classification of contigs

#contig	consensus taxid	consensus lineage
first1	210	cellular organisms; Bacteria; Proteobacteria; delta/epsilon subdivisions; Epsilonproteobacteria; Campylobacterales; Helicobacteraceae; Helicobacter; Helicobacter pylori
first10	1681	cellular organisms; Bacteria; Terrabacteria group; Actinobacteria; Actinobacteria; Bifidobacteriales; Bifidobacteriaceae; Bifidobacterium; Bifidobacterium bifidum
first100	1358	cellular organisms; Bacteria; Terrabacteria group; Firmicutes; Bacilli; Lactobacillales; Streptococcaceae; Lactococcus; Lactococcus lactis
first1000	1322	cellular organisms; Bacteria; Terrabacteria group; Firmicutes; Clostridia; Clostridiales; Lachnospiraceae; Blautia; Blautia hansenii
first10004	644	cellular organisms; Bacteria; Proteobacteria; Gammaproteobacteria; Aeromonadales; Aeromonadaceae; Aeromonas; Aeromonas hydrophila




Quantification table of reads on genes

id_cluster	first.featureCounts.tsv	second.featureCounts.tsv
first59.Prot_28193		844
first82.Prot_34066		847
first992.Prot_96159		4092
first8.Prot_06503		5279
first21.Prot_13879		584
		611

metagWGS will be used to analyze ExpoMycoPig data

How can we characterize microbiome digestive ecosystems for pig exposed to mycotoxins?

Experimental strategy



References

- [1] Di Tommaso et al. Nextflow enables reproducible computational workflows. *Nat Biotechnol.*, 2017.
- [2] Kurtzer et al. Singularity: Scientific containers for mobility of compute. *PLoS One*, 2017.
- [3] Kieser et al., ATLAS: a Snakemake workflow for assembly, annotation, and genomic binning of metagenome sequence data, bioRxiv, 2019.
- [4] Köster et al., Snakemake - A scalable bioinformatics workflow engine. *Bioinformatics*, 2012.
- [5] Martin. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal*, 2011.
- [6] Joshi and Fass. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files [Software]. Available at <https://github.com/najoshi/sickle>, 2011.
- [7] Li and Durbin. Fast and accurate short read alignment with Burrows-Wheeler Transform. *Bioinformatics*, 2009.
- [8] Menzel et al. Fast and sensitive taxonomic classification for metagenomics with Kaiju. *Nat Commun.*, 2016.
- [9] Ondov et al. Interactive metagenomic visualization in a Web browser. *BMC Bioinformatics*, 2011.
- [10] Andrews. FastQC. Available at <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>, 2010.
- [11] Nurk et al. MetaSPAdes: a new versatile metagenomic assembler. *Genome Res.*, 2017.
- [12] Li et al. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics*, 2015.
- [13] Li et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, 2009.
- [14] Seemann. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*, 2014.
- [15] Fu et al. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics*, 2012.
- [16] Liao et al. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, 2014.
- [17] Buchfink et al. Fast and sensitive protein alignment using DIAMOND. *Nature Methods*, 2015.
- [18] Langmead and Salzberg. Fast gapped-read alignment with Bowtie 2. *Nature Methods*, 2012.
- [19] Kang et al. MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ*, 2019.
- [20] von Meijenfeldt et al. Robust taxonomic classification of uncharted microbial sequences and bins with CAT and BAT. *Genome Biology*, 2019.
- [21] Ewels et al. MultiQC: Summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*, 2016.
- [22] Pinton and Oswald. Effect of deoxynivalenol and other Type B trichothecenes on the intestine: a review. *Toxins*, 2014.